Astronomical and Astrophysical Transactions (*AApTr*), 2012, Vol. 27, Issue 2, pp. 285–288,
ISSN 1055-6796, Photocopying permitted by license only, ©Cambridge Scientific Publishers

A simple model for tidal galactic interaction

I.A. Rezyapkin, L.P. Ossipkov*

Department of Space Technologies and Applied Astrodynamics, St. Petersburg State University, Stary Peterhof, St. Petersburg 198504 Russia

Received 1 February, 2012

Tidal interaction of galaxies is studied numerically on the basis of the restricted three-body problem. The evolution of an ensemble of test particles is considered for various cases of the relative motion of two point masses and for various values of the mass ratio.

Keywords: Celestial mechanics; Stellar dynamics; Methods: numerical; Galaxies: interaction

1 Introduction

Tidal interaction of galaxies is one of main factors governing their dynamical evolution. It was studied by many authors after pioneering works [1–6]. A deep analysis of the problem was fulfilled by A. & J. Toomre [7]. Note that authors [1–7] considered the restricted three-body problem.

The most of current studies of galaxy interaction and merging are based on N-body simulations (see e. g. a recent paper by Tutukov *et al.* [8] and a manual [9]). Many scenarios of galaxy evolution has appeared in course of discussions of obtained results. For instance, Byrd & Valtonen [10] put forward a hypothesis that SO galaxies result from spirals after destroying their spiral structure under the external gravitation of passing by galaxies.

But we think that some features of galaxy interaction can be revealed in frames of the restricted three-body problem. This model was used by some authors (e. g. [11–13]).

In this work we study an evolution of an ensemble of test points in a field of two gravitating point masses for various initial conditions. We shall restrict ourselves with the plane problem.

2 The equations of motion

Let m_A, m_B be masses of gravitating points A, B, X_B, Y_B be coordinates of the point B relatively the point A, and X, Y be relative coordinates of the test point.

*Corresponding author. Email: leonidosipkow@yahoo.com

The equations of motion will be as follows:

$$\ddot{X}+\frac{Gm_A}{R^3}X=Gm_B\left(\frac{X_B-X}{\Delta^3}-\frac{X}{R^3}\right),\qquad \ddot{Y}+\frac{Gm_A}{R^3}Y=Gm_B\left(\frac{Y_B-Y}{\Delta^3}-\frac{Y}{R^3}\right).$$

Here G is the gravitational constant, $R=\sqrt{X^2+Y^2}$, $\Delta=\left[(X_B-X)^2+(Y_B-Y)^2\right]^{1/2}$, $X_B=X_B(t)$ and $Y_B=Y_B(t)$ are considered as known functions.

Now it is necessary to transform the equations into a dimensionless form. We shall choose m_A as the mass unit, the least distance between points A and B, r_0, as the distance unit, $t_0=\left[r_0^3/(Gm_A)\right]^{1/2}$ as the time unit.

Denote

$$\mu=\frac{m_B}{m_A},\qquad x_B=\frac{X_B}{r_0},\ y_B=\frac{Y_B}{r_0},\qquad x=\frac{X}{r_0},\ y=\frac{Y}{r_0},\qquad \delta=\frac{\Delta}{r_0},\qquad \tau=\frac{t}{t_0}.$$

Then

$$\frac{d^2x}{d\tau^2}+\frac{x}{r^3}=\mu\left(\frac{x_B-x}{\delta^3}-\frac{x}{r^3}\right),\qquad \frac{d^2y}{d\tau^2}+\frac{y}{r^3}=\mu\left(\frac{y_B-y}{\delta^3}-\frac{y}{r^3}\right),$$

with $r^2=x^2+y^2$. These equations were integrated numerically by the Runge–Kutta method of the fourth order for $\tau\in[-5,10]$ under condition that $x_B^2+y_B^2=1$ for $\tau=1$.

3 Some results

We considered cases when the relative motion of the point B is parabolic, hyperbolic and elliptic. Below we shall discuss results for the parabolic orbit of point P.

At first we considered the simplest model when initial orbits of test points were circular. When the motion is direct, a transient spiral structure is formed and disappeared. There is no spiral pattern in the case of retrograde motions for $\mu=0.1$, but a ring structure can be formed from retrograde particles for larger μ.

When masses are equal ($\mu=1$), the most particles on direct orbits are captured by the passing point B. No spiral structure is formed, but tails are formed by thrown particles. Most of particles on retrograde orbits leave the system, and the rest ones form the ring structure.

When the orbits of test particles are elliptical for $\mu=0.5$ the influence of the passing point is not so significant.

In the case of the circular restricted three-body problem the resonance phenomena described by Lindblad [1] are significant for small μ. The cloud of test particles is deformed, and the spiral structure is developed. The larger μ, the more significant is the spiral structure that is transient and unstable in the case of direct orbits. The stable spiral structure develops in the case of retrograde orbits.

When the point B moves along a hyperbolic orbit, a relatively stable spiral structure results in the case of direct orbits of test particles.

Acknowledgements

The work of LO was supported in part by the Russian Foundation for Basic Research under grant 08–02–00361 and by a grant of President of the Russian Federation for support of Leading Scientific Schools NSh 3290.2010.2 "Kinematic and dynamic astronomy".

References

[1] P.O. Lindblad. Stockholms Observ. Ann. **20** 3 (1960).
[2] J. Pfleiderer and H. Siedentopf. Zs. Astrophys. **51** 201 (1961).
[3] J. Pfleiderer. Zs. Astrophys. **56** 12 (1964).
[4] N. Tashpulatov. Astron. Zh. **46** 1236 (1969).
[5] N. Tashpulatov. Astron. Zh. **47** 277 (1969).
[6] T.M. Eneev, N.N. Kozlov, and R.A. Sunyaev. Astron. Astrophys. **22** 41 (1973).
[7] A. Toomre and J. Toomre. Astrophys. J. **178** 623 (1972).
[8] A.V. Tutukov, V.V. Dryomov and G.N. Dryomova. Astron. Zh. **84** 487 (2007).
[9] V.V. Orlov and A.V. Rubinov. *N-Body Problem in Stellar Dynamics* (VVM Publ. House, SPb, 2008).
[10] G. Byrd and M. Valtonen. Astrophys. J. **350** 89 (1990).
[11] N.Ya. Sotnikova. Astrofizika **28** 495 (1980).
[12] V.N. Belov. Vestn. Leningrad. Univ., ser. 1. Iss. 4 105 (1990).
[13] A.V. Tutukov and A.V. Fedorova. Astron. Zh. **83** 589 (2006).